

100% SOLUBLE CALCIUM

Calcium Thiosulfate

0-0-0-10S-6Ca

LIQUID FERTILIZER

CaTs® – Calcium Thiosulfate Solution

Soluble Calcium (Ca)	6%
Combined Sulfur (S)	10%
Density: pound per gallon at 68°F	10.4
Volume: gallons per ton	192

GENERAL INFORMATION

CaTs® is a neutral to basic, chloride-free, clear solution, containing 6% calcium and 10% thiosulfate sulfur. Each gallon of CaTs contains 0.63 pound of calcium (Ca) and 1.0 pound of thiosulfate sulfur (S). CaTs may be applied by drip, sprinkler, or flood irrigation. It may be blended with other fertilizers or applied as a foliar treatment on selected crops. When used as a foliar fertilizer, **CaTs should first be diluted with water before applying. Blends of CaTs should not be acidified below a pH of 6.0.**

CaTs may be used as a fertilizer for the correction of calcium and sulfur deficiency as well as a soil amendment. As a soil amendment CaTs may be used to improve water infiltration and aid in leaching of harmful soil salts.

CaTs is compatible with most fertilizer solutions.

CaTs is not compatible with phosphate, sulfate and ammonium thiosulfate fertilizers. In absence of specific recommendations and data, do a jar test before mixing large quantities. The addition of water to the mixture may be helpful to maintain blend stability.

When blending with micronutrients and pesticides, trial blends should be conducted before beginning large scale mixing. When mixing other liquid fertilizers and/or pesticides with CaTs, the blend sequence should be as follows: water, pesticide, CaTs and/or other fertilizer.



THE 100% SOLUBLE CALCIUM SOIL AMENDMENT

Delivers Maximum Efficiency

APPLICATION & USE RECOMMENDATIONS

See Application Precautions
on Page 3 Before Applying

CaTs may be applied to a wide variety of ornamental, turf, greenhouse, and other agricultural crops. Calcium requirement for most crops increases during periods of rapid growth and early fruit development. Application of CaTs should be made based on soil, fruit and/or plant tissue analysis for calcium.

SOIL APPLICATION

CaTs may be broadcast, banded or knife injected as a source of calcium and sulfur. Rates may vary between 20 to 40 gallons per acre. CaTs is not compatible with phosphate fertilizers. Preplant treatment, apply 30 to 40 gallons of CaTs per acre via irrigation water depending on the salt content in the soil that requires leaching. CaTs can be used in a program to reduce problems related to calcium disorders in tree fruit. Soil application of CaTs should be used (and is more effective) in conjunction with a foliar calcium program.

Apples, Pears, Cherries, Apricots, Filberts, Peaches: Producing trees (at least 5 years old) on sandy loam soil – apply 20 to 30 gallons per acre in a 4 foot band under the drip line. Irrigate after application.

Young Trees and/or Trees on Sandy or Light Soils: Apply 15 to 20 gallons per acre. Irrigate after application.

Example: A 10 acre orchard at 20 gallons per acre would require 200 gallons applied in a 4 foot band.

SOIL AMENDMENT

CaTs may be broadcast or band applied to the soil surface to reduce crusting problems and improve water infiltration. For band application, apply at the rate of 1 gallon per acre per inch of band width. (An 8 inch band would require an 8 gallon per acre rate.) Preplant treatment, apply 20 to 40 gallons of CaTs per acre via irrigation water. CaTs may be used as a soil amendment during the growing season to reduce the harmful effects of sodium. CaTs may be applied with irrigation water or directly to the soil before irrigation. Rate of application is dependent on severity of the sodium problem. Contact a Tessenderlo Kerley, Inc. representative for additional information.

IRRIGATION WATER TREATMENT

Irrigation water derived from snowmelt or rainwater may contain low levels of salts (low electrical conductivity). While this is good for crops, it can have a negative effect on water infiltration in many soils, resulting in puddling and/or runoff. Crops become moisture stressed during periods of high temperatures due to lack of deep moisture and/or shallow root system.

Application of CaTs can:

1. improve water infiltration
2. increase deep moisture and
3. reduce moisture stress.

Application rates are dependent on field conditions. For additional information contact a Tessenderlo Kerley representative.

FERTIGATION

CaTs may be applied through most irrigation systems. Apply CaTs in the middle 1/3 of the irrigation set.

OVERHEAD IRRIGATION

Apples: Apply 5 to 10 gallons per acre per application. Repeat as needed to achieve calcium tissue levels desired.

Young Tree and Vine Crops: Apply 3 to 8 gallons per acre per application. Repeat as required.

Mature Tree and Vine Crops: Apply 5 to 10 gallons per acre per application. Repeat as required.

Potatoes: Apply 10 to 15 gallons per acre per application beginning at hooking stage. Apply a second application when the potatoes are about golf ball size.

Onions: Apply 10 to 12 gallons per acre beginning after third leaf stage. Apply 2 to 3 times with at least 1 week intervals.

Alfalfa: Apply 10 gallons per acre applied after the first cutting and again after the third cutting.

Other crops: Apply 5 to 10 gallons per acre per application. Repeat as required.

Ornamentals: Apply 1 quart of CaTs per 1000 square feet in irrigation water to growing plants. Rinse off any contact with foliage immediately.

Turf: Apply CaTs in irrigation water at 30 to 40 ounces per 1000 square feet. Apply as needed during the growing season as indicated by soil and tissue testing.

FLOOD AND FURROW

Row Crops: Apply 10 to 15 gallons per acre per application with irrigation water to improve water infiltration and to reduce runoff. Apply as needed during the growing season.

Tree Crops: Apply 10 to 15 gallons per acre per application with irrigation water to improve water infiltration and to reduce runoff. Apply as needed during the growing season.

FOLIAR APPLICATION

Apply in sufficient volume to achieve good plant coverage, usually 20 to 30 gallons per acre for row crops.

FIELD CROPS

Alfalfa: Apply 2 to 4 gallons per acre per application. Apply on stubble of previous cutting.

Dry Peas: Apply 1 to 1.5 gallon per acre per application at 1/10th bloom stage.

Lentils: Apply 1 to 1.5 gallon per acre per application at 1/10th bloom stage.

Onions: Apply 6 to 8 gallons per acre after the 3rd leaf stage when crop is well established. Apply 2 to 4 times during the season.

Other Crops: Apply 1/2 to 1 gallon per acre per application. Repeat applications as required.

TREE CROPS

For calcium related disorders, begin applications starting at petal fall.

Apples (Gala and Red Delicious): Apply 2 to 3 quarts/100 gallons per acre. Apply 4 to 6 times during the growing season.

Pears (Bartlett, D'Anjou, Bosc, Concord and Arirang Asian): Apply 2 to 4 quarts/100 gallons per acre. Begin application at petal fall and continue through the season for 4 to 6 times.

Cherries: Apply 2 to 3 quarts/100 gallons per acre beginning at petal fall. Apply every 7 to 10 days up to four times or at the beginning and during a rainfall to reduce cracking. **DO NOT apply CaTs with a crop oil spray. Allow at least 5 days before or 9 days after an oil spray before applying CaTs as a foliar.**

Combination of other fertilizers and pesticides with CaTs may increase the chances of foliar burn. It is beyond the capability of Tessengerlo Kerley, Inc. to test all combinations of foliar fertilizers and/or pesticides. The dealer/grower should conduct a test plot when applying with multiple products before beginning large scale applications. When mixing with unfamiliar products, always do a jar test to check for compatibility.

DRIP IRRIGATION

Young Tree and Vine Crops: Apply 4 to 8 gallons per acre per application during the growing season. Allow 10 to 14 days between applications.

Mature Trees and Vine Crops: Apply 5 to 10 gallons per acre per application. Repeat two to three times during the growing season or as needed. Allow 7 to 8 days between applications.

Vegetable and Row Crops: Apply 3 to 5 gallons per acre per application. Repeat 4 to 5 times during the growing season as needed.

APPLICATION PRECAUTIONS

CaTs is not compatible with phosphate fertilizers.

DO NOT apply CaTs to foliage of crops sensitive (foliar burn) to sulfur such as corn.

DO NOT mix with acids or acidic fertilizers below a pH of 6.0. **Be sure to check other manufacturers' labels concerning dormant oil treatment spray guidelines and foliar nutrient applications containing sulfur.**

DO NOT apply CaTs foliar with crop oil sprays. Allow at least 5 days before or 9 days after an application of crop oil before applying CaTs as a foliar.

DO NOT apply CaTs or other thiosulfate fertilizers while chlorinating irrigation system. Thiosulfates will neutralize chlorine.

DO NOT apply to foliage of any crop when temperatures are above 90°F. Apply CaTs in the early morning or late evening.

When mixing CaTs or any liquid fertilizer with pesticides, always keep agitators running during filling and spraying operations. Failure to maintain agitation may cause separation of products resulting in uneven spray application.

Pop Up Fertilizers: Many crops are sensitive to salts during germination. When soil moisture is low, delayed crop emergence and/or phytotoxicity may occur when fertilizer is placed too close to the seed. **DO NOT** use CaTs in pop up fertilizer when soil moisture is limited; soil salinity is above an electrical conductivity of 2; or when irrigation is delayed such that germination may be affected. **DO NOT apply CaTs with knife injectors or other types of fertilizer injecting equipment that may cause root pruning.**

Sprinkler application of CaTs and other liquid fertilizers over an established crop may cause foliar injury to a crop if: injection period is short enough to cause an excessive amount of fertilizer to accumulate on the leaves; temperatures are above 90°F and humidity less than 30%; fertilizer rates are higher than recommended; irrigation pump breaks down during or immediately after injecting fertilizer; and/or any combination of these conditions.

PHYTOTOXICITY

Plant and leaf injury may occur on some crops when certain weather and growing conditions are present. The user assumes all risks of use and handling.

EXAMPLES OF BLENDS WITH CaTs

PRODUCT	LBS / TON	ANALYSIS
CaTs	1,600	0-0-9-8S-4.8Ca
KCl (Dry 62%)	290	
Water	110	
CaTs	1,000	4.5-0-0-5S-8.5Ca
CN-9	1,000	
CaTs	1,667	7-0-0-8S-5Ca
Urea (Dry)	304	
Water	29	

Keep out of reach of children. Use caution when handling.

See SDS for additional information on safety and handling at: cropvitality.com/cats



Technical Data

CaTs® 0-0-0-10S-6Ca

PLANT NUTRIENT CONTENT WEIGHT %

Calcium (Ca)	6
Total Sulfur (S)	10

TYPICAL PROPERTIES

Specific Gravity	1.25
pH	6.5 - 8.0
Appearance	Colorless
Salt-Out Temperature	32°F

FORMULATION AND APPLICATION FACTORS, 68°F

Density: pounds per gallon	10.4
Volume: gallons per ton	192
Pounds of Calcium per gallon	0.63
Pounds of Sulfur per gallon	1.0

The manufacturing process of CaTs is patent protected.

Warranty and Limitation of Damages

Tessenderlo Kerley, Inc. (TKI) warrants only that this product conforms to the product description in the Application Guide. Except as warranted by this description, TKI makes no representation or warranty or guarantee, whether expressed or implied, of fitness for a particular purpose of merchantability, or of product performance. TKI does not authorize any agent or representative to make any such representation, warranty or guarantee. To the extent consistent with applicable law, TKI's maximum liability for breach of its warranty or for use of this product, regardless of the form of action, shall be limited to the purchase price of this product. To the extent consistent with applicable law, buyer and user acknowledge and assume all risks and disposal liability resulting from handling, storage, use and disposal of this product. If buyer does not agree with or accept these warranty and liability limitations, buyer may return the unopened container to the place of purchase for full refund. Buyer's use of this product shall constitute conclusive evidence of buyer's acknowledgement and acceptance of the forgoing limitations. Some jurisdictions do not allow the exclusion of implied warranties or the limitation of certain damages, so the above may not apply. The purchase, delivery, acceptance and use of this product by the buyer are subject to the terms and conditions of seller's sales invoice for this product.

©2003, 2010, 2013 Tessenderlo Kerley, Inc. All rights reserved. This material is protected by the copyright laws of the United States and international treaties. Use, copying, reproduction, distribution, display or transmission in any form by any means, without the express written permission of Tessenderlo Kerley, Inc. is prohibited and may result in civil and criminal liability.

Contact Information

Tessenderlo Kerley, Inc.
2255 North 44th Street, Suite 300 | Phoenix, AZ 85008
Telephone: 602-889-8300 | Toll-Free: 800-525-2803
email: cats@cropvitality.com | cropvitality.com

CaTs® is a registered trademark of Tessenderlo Kerley, Inc.



Revised: 07/2013